

ROUTINE GLUCOMMANDER ORDERS

Admit as an inpatient to the services of Dr. _____

Notify The Diabetes Treatment Center of insulin drip order via Meditech

Diagnosis: _____

IF PATIENT DIAGNOSED WITH DKA OR HHNK REFER TO APPROPRIATE DKA OR HHNK ORDERS

Parameters for Glucomander Set -up	a. Low target blood glucose= 100mg/dl b. High target blood glucose = 140 mg/dl c. Maximum minutes for checking blood glucose = 60 d. Initial multiplier = 0.02
Dietary	Diet: Non-caloric unless NPO or Continuous tube feedings of _____ at _____ ml/hour
Laboratory	1. CHEM BASIC 2. Phosphorus 3. Calcium 4. CBC
Medications	1. IV Fluids: Normal Saline 1000 ml at 21 ml/hr, when blood glucose <250, change to D5W/0.9% NaCl at the same rate. Other IV Fluid: _____ _____ 2. Insulin Drip: Regular Insulin 250 units added to 0.9% NaCl 250 ml
Criteria for stopping drip	2. When the blood glucose is between 100-140 X4, and C02 >18 or anion gap < 14 a. Contact MD for order to discontinue drips, start diet, home diabetes medications and sliding scale insulin b. Once home medications are resumed, wait one hour before stopping insulin drip.

Physician Signature: _____ Date / Time: _____



Routine Glucomander Orders